Astrophysics and Space Science manuscript No.
(will be inserted by the editor)

arXiv:astro-ph/0608311v1 15 Aug 2006

Bryan M. Gaensler · Maura McLaughlin · Stephen Reynolds · Kazik Borkowski · Nanda Rea · Andrea Possenti · Gianluca Israel · Marta Burgay · Fernando Camilo · Shami Chatterjee · Michael Kramer · Andrew Lyne · Ingrid Stairs

Chandra Smells a RRAT

X-ray Detection of a Rotating Radio Transient

Received: date / Accepted: date

Abstract "Rotating RAdio Transients" (RRATs) are a newly discovered astronomical phenomenon, characterised by occasional brief radio bursts, with average intervals between bursts ranging from minutes to hours. The burst spacings allow identification of periodicities, which fall in the range 0.4 to 7 seconds. The RRATs thus seem to be rotating neutron stars, albeit with properties very different from the rest of the population. We here present the serendipitous detection with the *Chandra X-ray Observatory* of a bright point-like X-ray source coincident with one of the RRATs. We discuss the temporal and spectral properties of this X-ray emission, consider counterparts in other wavebands, and interpret these results in the context of possible explanations for the RRAT population.

Keywords pulsars: individual (J1819–1458) · stars: flare, neutron · X-rays: stars

PACS 97.60.Gb · 97.60.Jd · 98.70.Qy

B. M. Gaensler and S. Chatterjee
Harvard-Smithsonian Center for Astrophysics, Cambridge MA, USA
E-mail: bgaensler@cfa.harvard.edu
Present address: The University of Sydney, NSW, Australia

M. McLaughlin
University of West Virginia

S. Reynolds and K. Borkowski
North Carolina State University

N. Rea
SRON Netherlands Institute for Space Research

A. Possenti and M. Burgay
Osservatorio Astronomico di Cagliari

G. Israel
Osservatorio Astronomico di Roma

F. Camilo
Columbia University

M. Kramer and A. Lyne
Jodrell Bank Observatory

I. Stairs
University of British Columbia

1 Introduction

Astronomers are still coming to terms with the fact that the zoo of isolated neutron stars harbours an increasingly diverse population. Objects of considerable interest include radio pulsars, anomalous X-ray pulsars, soft gamma repeaters, central compact objects in supernova remnants (SNRs), and dim isolated neutron stars. A startling new discovery has now forced us to further expand this group: McLaughlin et al. (2006) have recently reported the detection of eleven "Rotation RAdio Transients", or "RRATs", characterised by repeated, irregular radio bursts, with burst durations of 2–30 ms, and intervals between bursts of ~ 4 min to ~ 3 hr. The RRATs are concentrated at low Galactic latitudes, with distances implied by their dispersion measures of $\sim 2-7$ kpc.

For ten of the eleven RRATs discovered by McLaughlin et al. (2006), an analysis of the spacings between repeat bursts reveals an underlying spin period, P, and also in three cases, a spin period derivative, $\dot{P}$. The observed periods fall in the range $0.4 \text{ s} < P < 7$ s, which generally overlap with those seen for the radio pulsar population.

For the three RRATs with values measured for both P and $\dot{P}$, a characteristic age, τ_c, and a dipole surface magnetic field, B, can both be inferred, as listed in Table 1. These sources can also be placed on the standard "$P-\dot{P}$ diagram", and can thus be compared to other populations of rotating neutron star:

- RRAT J1317–5759 has properties typical of radio pulsars, except for a relatively long spin period.
- RRAT J1819–1458 is very young, with a high magnetic field. On the $P-\dot{P}$ diagram, it is located in the upper-right region, in the same area occupied by the magnetars and by the high-field radio pulsars.
- RRAT J1913+3333 has spin properties indistinguishable from the bulk of radio pulsars.

We here present the X-ray detection of RRAT J1819–1458, which provides a further point of comparison with the various neutron star population. The details of this detection and its interpretation are discussed by Reynolds et al. (2006).

Table 1 Properties of the three RRATs with measured period derivatives.

Source	P (sec)	τ_c (Myr)	B (10^{12} G)
RRAT J1317–5759	2.6	3.3	5.83
RRAT J1819–1458	4.3	0.12	50
RRAT J1913+3333	0.92	1.9	2.7

2 X-ray Emission from RRAT J1819–1458

2.1 Detection

As mentioned above, RRAT J1819–1458 is very young, and has a high surface magnetic field. This source bursts every ~ 3 min, making it the most active of the known RRATs. Its dispersion measure of 196 ± 3 pc cm^{-3} corresponds to an estimated distance of 3.6 kpc, with considerable uncertainty.

RRAT J1819–1458 sits only $\sim 11'$ from the young SNR G15.9+0.2. This source was the target of a 30 ks observation with *Chandra* ACIS in May 2005, and RRAT J1819–1458 fortuitously falls within the field of view (Reynolds et al., 2006). As shown in Figure 1, there is a clear detection of a bright, unresolved X-ray source within the error ellipse for RRAT J1819–1458. Using the X-ray differential source count distribution for the Galactic plane of Sugizaki et al. (2001), we find that the probability of finding a field source of this count rate at this position is $< 10^{-4}$, and conclude that we have both identified and localised the X-ray counterpart of RRAT J1819–1458.

2.2 Spectrum and Variability

We have extracted 524 ± 24 counts from the X-ray counterpart to RRAT J1819–1458, the spectrum from which is shown in Figure 2. While this is insufficient for detailed spectral modelling, it still yields crucial information.

Fitting simple absorbed power-law and blackbody models, we find that the spectrum is a poor fit to the former, but a good fit to the latter (Reynolds et al., 2006). At a distance of $3.6d_{3.6}$ kpc, the inferred blackbody radius (as viewed at infinity) is $20d_{3.6}$ km, which is consistent with standard neutron star equations of state, given the uncertainty in the distance estimate. For this blackbody fit, the foreground absorbing column is $N_H = 7^{+7}_{-4} \times 10^{21}$ cm^{-2}, and the surface temperature (at infinity) is $kT_\infty = 120 \pm 40$ eV. In the energy range 0.5–8.0 keV, the unabsorbed flux is 2×10^{-12} ergs cm^{-2} s^{-1} and the isotropic luminosity is $3.6d_{3.6}^2 \times 10^{33}$ ergs s^{-1}.



Fig. 1 A *Chandra* ACIS image of RRAT J1819–1458, observed $11'$ off-axis, and smoothed with a gaussian of FWHM $2''$. The ellipse shows the $3 - \sigma$ error ellipse for the position of the RRAT, as derived by McLaughlin et al. (2006) from radio timing.



Fig. 2 The X-ray spectrum of RRAT J1819–1458, as observed with *Chandra* ACIS (Reynolds et al., 2006). The data points show the measurements, while the solid line shows the best-fit absorbed blackbody described in the text.

Given the extreme variability of the source seen at radio wavelengths, it is of interest to quantify the level of X-ray variability seen in this observation. Examination of individual CCD frames (at a time-resolution of 3.2-seconds) shows no evidence for individual brief X-ray bursts that might be a counterpart to the RRAT phenomenon, with a 3-σ upper limit on the observed fluence of any burst of 3×10^{-11} ergs cm^{-2} in the 0.5–8 keV energy range. The data also show no evidence of variability on any time-scale ranging from 3.2 s to 5 days (the time-span covered by the observations). Although the time resolution is only slightly shorter than the spin period,



Fig. 3 A 2MASS image of the field surrounding RRAT J1819–1458, with J, H and K bands shown in blue, green and red, respectively. The arrows mark the position of the RRAT, as derived from *Chandra* observations. The upper limits on the infrared emission from RRAT J1819–1458 are $J > 15.6$, $H > 15.0$ and $K > 14.0$.

$P = 4.3$ s, we do have limited sensitivity to an aliased pulsed signal. However, we find no pulsations down to a 3-σ pulsed fraction limit of 70% for a sinusoidal pulse profile (Reynolds et al., 2006).

2.3 Data At Other Wavelengths

With an accurate position for RRAT J1819–1458 in hand from our *Chandra* position, we have searched for a counterpart to this source at other wavelengths, using archival VLA, 2MASS and GLIMPSE data. No detection is made in any of these data.

The non-detection in 2MASS data, shown in Figure 3, allows us to put a lower limit on the X-ray to infrared flux ratio of $f_X/f_{IR} > 0.7$, which rules out most stellar counterparts (see Kaplan et al., 2004). Furthermore, the foreground column density inferred from the X-ray spectral fit is substantially smaller than the integrated column through the Galaxy at this position, arguing against the X-ray emission being from a background galaxy or cluster. These arguments, combined with the good blackbody fit to the X-ray spectrum, imply that the most reasonable interpretation for the X-ray emission is thermal emission from a neutron star surface.

3 Comparison with Other Sources

It is unclear whether the RRATs are a completely new group of neutron stars, or are a new manifestation of one of the previously known classes of object. Since all classes of young neutron star are X-ray emitters, the detection reported here provides vital new information which allows us to begin to discriminate between various interpretations for the RRATs.

We first note that while many neutron stars are pulsed in X-rays, those which produce thermal emission inevitably exhibit low pulsed fractions, due to gravitational bending of the light emanating from their surfaces (e.g., Psaltis et al., 2000). Thus our relatively poor limit on pulsed fraction is unsurprising and unconstraining.

Despite its location near the magnetars on the $P - \dot{P}$ diagram, the X-ray properties of RRAT J1819–1458 are distinct from those seen for magnetars: the RRAT is much colder and less luminous than the magnetars, and apparently lacks the hard X-ray tail seen for these sources, at our current level of sensitivity. Furthermore, the magnetar birth rate is well below that estimated for the RRAT population (Popov et al., 2006). The only possible link to magnetars is with the transient magnetar XTE J1810–197 which, while in quiescence, had a surface temperature $kT_\infty \approx 150 - 180$ eV (Ibrahim et al., 2004; Gotthelf et al., 2004), comparable to that seen here.

The X-ray temperature of RRAT J1819–1458 is also well below those of the "central compact objects", while its temperature and luminosity are above those of most of the dim isolated neutron stars.

The one population whose X-ray properties provide a reasonable match to that of the RRAT is that of radio pulsars, for which sources with ages around 100 kyr show spectra very similar to what we have found here. For example, PSR J0538+2817 is 30 kyr old and has $kT_\infty = 160$ eV, while PSR B0656+14 is 110 kyr old and has $kT_\infty = 70$ eV (see Reynolds et al., 2006, for details). The X-ray emission from RRAT J1819–1458 thus suggests that this source is a normal radio pulsar, albeit one that produces unusual radio bursts. Additional evidence to support this possibility is the subsequent discovery from PSR B0656+14 of RRAT-like behaviour (Weltevrede et al., 2006). If PSR B0656+14 was placed at the distance of RRAT J1819–1458, its occasional bright radio bursts would still be seen, but not the underlying regular train of pulsations.

One point to note is that the inferred surface magnetic field strength of RRAT J1819–1458 is more than an order of magnitude greater than those of PSRs J0538+2817 and B0656+14 discussed above. Two radio pulsars with comparable magnetic fields that have been detected in X-rays are PSRs J1718–3718 (Kaspi and McLaughlin, 2005) and J1119–6127 (Gonzalez et al., 2005). These sources show temperatures ($kT \sim 150-200$ eV) and luminosities ($\sim 10^{32}$–10^{33} ergs s^{-1}) comparable to that of RRAT J1819–1458, although both sources are probably much younger

(35 and 1.7 kyr, respectively) and, in contrast to RRAT J1819–1456, have X-ray luminosities less than their spin-down luminosities.

4 Possible Interpretations

If the RRATs are normal radio pulsars as proposed above, we then need to explain their transient behaviour. One possibility, discussed by Zhang et al. (2006), is that RRATs are pulsars that are no longer active, but for which a temporary "star spot" with multipole field components emerges above the surface. This magnetic field component could temporarily reactivate the radio beaming mechanism, producing the observed bursts.

The difficulty with this possibility is that none of the RRATs with known values of $\dot{P}$ appear to be near the "death line" in the $P - \dot{P}$ diagram, beyond which the pulsar mechanism is expected to turn off. To account for this, Zhang et al. (2006) have proposed that RRATs have dipole fields offset from their centres, causing their magnetic fields to be over-estimated. However, the X-ray temperature seen for RRAT J1819–1458 is consistent with it being a $\sim$ 100-kyr old neutron star (see Sec. 3 above), in reasonable agreement with its characteristic age as listed in Table 1, and confirming that if this is a radio pulsar, it is as yet nowhere near death.

Zhang et al. (2006) also consider the possibility that RRATs are caused by a brief reversal of the direction in which radio beams are emitted. They draw upon the previous work of Gil et al. (1994), who showed that whenever PSR B1822–09 produces "nulls" in its main pulse, it produces a much weaker interpulse, which has been interpreted by Dyks et al. (2005) to correspond to radiation temporarily emitted in the opposite direction. If one invokes an emitting geometry in which the main pulse is never seen, then the interpulse alone, appearing only when the unseen main pulse nulls, might correspond to RRAT-like emission. The difficulty with this interpretation is that in PSR B1822–09 this reversal lasts for several minutes, in contrast to the RRATs, for which multiple bursts in succession are yet to be observed.

Finally, the RRAT mechanism might be produced by interaction of the neutron star with an equatorial fallback disk or with orbiting circumpolar debris. Accretion from a disk should usually quench the radio emission mechanism, but sporadic drops in the accretion rate could allow the radio beam to turn on for a fraction of a second, producing the RRAT phenomenon (Li, 2006). However, this possibility is at odds with the behaviour seen by Weltevrede et al. (2006) for PSR B0656+14, in which the RRAT-like bursts are superimposed on an underlying persistent series of faint radio pulsations. Alternatively, episodic injection of material from a circumpolar asteroid belt could temporarily activate a quiescent region of the magnetosphere, producing a RRAT burst, in some cases from a radio pulsar that is normally beaming away from us (Cordes and Shannon, 2006).

5 Future Observations and Conclusions

A variety of forthcoming observations should be able to cast more light on the possibilities discussed above, and on the spatial and spin distributions of these sources. Several groups have already undertaken new radio searches for RRAT-like emission from other classes of neutron stars. Near-infrared observations with the VLT have also been obtained for some RRATs, to provide stronger constraints on the X-ray to infrared flux ratio of these sources, and to look for hints of a fossil / fallback disk (see Wang et al., 2006). Meanwhile, in a project called "Astro Pulse", the data from the Arecibo SETI@Home project are being reanalysed for short transient signals. With the sensitivity of Arecibo, many RRATs may be found in this analysis.

Finally, following on from our X-ray detection discussed here, *XMM-Newton* observations of two RRATs have been approved for Cycle 5: of RRAT J1819–1458 (to obtain a better spectrum and to properly search for pulsations) and of RRAT J1317–5759 (to see if it too emits detectable X-rays).

A final point to make is that it is already clear from the 11 RRATs known that the birth rate of these objects is $\sim 3 - 4$ times that previously estimated for all radio pulsars (McLaughlin et al., 2006; Popov et al., 2006). We are thus only seeing the upper tip of a large transient distribution. When one combines this with the populations of nullers (Backer, 1970), "bursters" (e.g., PSR J1752+2359; Lewandowski et al., 2004), "winkers" (PSR B1931+24; Kramer et al., 2006) and "burpers" (GCRT J1745-3009; Hyman et al., 2005), it becomes clear that a full study of the transient radio sky is needed (Cordes et al., 2004). The next generation of radio telescopes, with the capability of monitoring very wide fields of view (e.g., LOFAR, xNTD and the SKA) should make many further discoveries of such phenomena.

Acknowledgements B.M.G. acknowledges the support of NASA through LTSA grant NAG5-13023 and of an an Alfred P. Sloan Fellowship.

References

Backer D.C. Pulsar nulling phenomena. Nature, **228**, 42 (1970)

Cordes J.M. & Shannon, R.M. Rocking the lighthouse: Circumpulsar asteroids and radio intermittency. ApJ, submitted (astro-ph/0605145) (2006)

Cordes J.M., Lazio, T.J.W., & McLaughlin M.A.. The dynamic radio sky. New Astronomy Reviews, **48**, 1459 (2004)

Dyks J., Zhang B., & Gil, J. Reversals of radio emission direction in PSR B1822–09. ApJ, **626**, L45 (2005)

Gil J.A., Jessner A., Kijak J. et al. Multi-frequency study of PSR 1822−09. A&A, **282**, 45, (1994)

Gonzalez M.E., Kaspi V.M., Camilo F., et al. Unusual Pulsed X-Ray Emission from the Young, High Magnetic Field Pulsar PSR J1119-6127. ApJ, **630**, 489 (2005)

Gotthelf E.V., Halpern J.P., Buxton M., & Bailyn C. Imaging X-Ray, Optical, and Infrared Observations of the Transient Anomalous X-Ray Pulsar XTE J1810–197. ApJ, **605**, 368 (2004)

Hyman S.D., Lazio T.J.W., Kassim N.E., et al. A powerful bursting radio source towards the Galactic Centre. Nature, **434**, 50 (2005)

Ibrahim A.I., Markwardt C.B., Swank J.H., et al. Discovery of a Transient Magnetar: XTE J1810–197. ApJ, **609**, L21 (2004)

Kaplan D.L., Frail D.A., Gaensler B.M., et al. An X-Ray Search for Compact Central Sources in Supernova Remnants. I. SNRS G093.3+6.9, G315.4-2.3, G084.2+0.8, and G127.1+0.5. ApJS, **153**, 269 (2004)

Kaspi V.M., & McLaughlin M.A. Chandra X-ray Detection of the High-Magnetic-Field Radio Pulsar PSR J1718-3718. ApJ, **618**, L41 (2005)

Kramer M., Lyne A.G., O'Brien J.T., et al. A periodically active pulsar giving insight into magnetospheric physics. Science, **312**, 549 (2006)

Lewandowski W., Wolszczan A., Feiler G., et al. Arecibo timing and single-pulse observations of eighteen pulsars. ApJ, **600**, 905 (2004)

Li X.-D.. On the nature of part time radio pulsars. ApJ, **646**, L139 (2006)

McLaughlin M.A., Lyne A.G., Lorimer D.R., et al. Transient radio bursts from rotating neutron stars. Nature, **439**, 817 (2006)

Popov S.B., Turolla R., & Possenti, A. A tale of two populations: rotating radio transients and X-ray dim isolated neutron stars. MNRAS, **369**, L23 (2006)

Psaltis D., Özel F., & DeDeo S. Photon propagation around compact objects and the inferred properties of thermally emitting neutron stars. ApJ, **544**, 390 (2000)

Reynolds S., Borkowski K., Gaensler B., et al. Discovery of the X-ray counterpart to the rotating radio transient J1819–1458. ApJ, **639**, L71 (2006)

Sugizaki M., Mitsuda K., Kaneda H., et al. ApJS, **134**, 77 (2001)

Wang Z., Chakrabarty D., & Kaplan D.L. A debris disk around an isolated young neutron star. Nature, **440**, 772 (2006)

Weltevrede P., Stappers B.W., Rankin J.M., & Wright G.A.E. Is pulsar B0656+14 a very nearby RRAT source? ApJ, **645**, L149 (2006)

Zhang B., Gil J., & Dyks J. Physical interpretations of rotating radio transients. ApJ, submitted (astro-ph/0601063) (2006)